

**Supplementary Table 1.** Several reported cases of somnambulism associated with OSA in the literature

Case number	Demographics	Concomitant sleep disorder	Prior treatment	Successful treatment option
Current study	12/M	Moderate OSA (AHI: 14.1/h) sleepwalking/talking/terror	None	Auto-set CPAP (4–15 cmH <sub>2</sub> O) residual AHI: 1.7/h after treatment
Tale et al. <sup>1</sup>	42/M	Moderate OSA (AHI: 22.4/h) sleepwalking/incomplete sleep paralysis	CPAP (non-compliant)	Clonazepam 0.25 mg at bedtime
Sun et al. <sup>2</sup>	46/M	Severe OSA (AHI: 40.1/h) sleepwalking/terror	Tablet clonazepam 2 mg at bedtime (felt worse; denied taking medication)	Auto-set CPAP (5–18 cmH <sub>2</sub> O) residual AHI: 3.2/h after treatment
Soca et al. <sup>3</sup>	42/M	Severe OSA (AHI: 41/h) sleepwalking/talking/sex REM sleep behavior disorder	None	CPAP (10 cmH <sub>2</sub> O) improved sleepwalking/talking with clonazepam 0.5 mg at bedtime improved sleep and sex
Lateef et al. <sup>4</sup>	54/F	OSA sleepwalking/driving	None	CPAP

OSA, obstructive sleep apnea; AHI, apnea-hypopnea index; CPAP, continuous positive airway pressure

## REFERENCES

1. Tale S, Kumar A, Saini LK, Meitei SP, Gupta R. Sleepwalking and prolonged partial sleep paralysis in a case of obstructive sleep apnea. *Sleep Sci* 2021;14(2):186-189.
2. Sun Y, Li J, Zhang X, Jiao Q, Yang S, Ji L. Case report: parasomnia overlap disorder induced by obstructive sleep hypopnea apnea syndrome: a case report and literature review. *Front Neurosci* 2020;14:578171.
3. Soca R, Keenan JC, Schenck CH. Parasomnia overlap disorder with sexual behaviors during sleep in a patient with obstructive sleep apnea. *J Clin Sleep Med* 2016;12(8):1189-1191.
4. Lateef O, Wyatt J, Cartwright R. A case of violent non-REM parasomnias that resolved with treatment of obstructive sleep apnea. *Chest* 2005;128(4 Suppl):461S.